

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

WARNING: May be fatal if swallowed. Wash thoroughly with soap and water after handling and before eating or smoking. This product can be absorbed through the skin. Wear a respirator and gloves when applying or handling this product. Keep away from humans, domestic animals and pets. Any person who retrieves carcasses or unused bait following application of this product must wear gloves. All handlers (including applicators) must wear long sleeved shirt and long pants, shoes plus socks, gloves, protective eyewear and a dust/mist filtering respirator (NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

FIRST AID: Have this label with you when obtaining treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin with plenty of cool water for 15-20 minutes. Call a poison control center or doctor for treatment advice. If inhaled: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

TREATMENT FOR PET POISONING: If animal eats tracking powder, call veterinarian at once.

NOTE TO VETERINARIAN: Anticoagulant Chlorophacinone: For animals ingesting tracking powder, and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Store in original container in a cool, dry place inaccessible to children and pets.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Handling: This is a nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or dispose of empty container in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions

RESTRICTED USE PESTICIDE DUE TO HIGH ACUTE TOXICITY

For retail sale to and use by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certificate.



TRACKING POWDER

For Use Indoors
and in Burrows Leading Into Buildings Only

Active Ingredient: chlorophacinone 0.2%
Inert Ingredients 99.8%
Total 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING: See side panel for additional precautionary statements.

LIPHATECH®

Liphatech, Inc.
3600 W. Elm Street
Milwaukee, WI 53209
(800) 351-1476

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: Use only against Norway rats, Roof rats and House mice inside of homes, industrial and agricultural buildings, and similar man made structures. Rozol Tracking Powder may also be dusted into rat burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures. Tracking powder must be placed in locations not accessible to children, pets, domestic animals or non-target wildlife. Do not apply powder in areas where there is a possibility of contaminating water, food, feedstuffs, food or food handling equipment, or milk or meat handling equipment or surfaces that come in direct contact with food. Applications in these areas are limited only to concealed, inaccessible places such as wall voids. Do not apply powder along walls or in corners or in open floor areas. Do not place near or inside ventilation duct openings. Use only for sites, pests, and application methods described on this label.

SELECTION OF TREATMENT AREAS: Determine dry areas where rats and house-mice will most likely pick up the powder on their feet or fur and ingest it during grooming. Generally, these areas are along walls by gnawed openings and burrows, in corners and concealed places, in spaces between floors and walls, or in locations where rodents or their signs have been observed. Remove goods piled directly on the floor and place on skids. Use boxes or other obstacles to force rodents to travel through constricted areas. Give special attention to the climbing ability of Roof rats. For this species, use suitable PVC, sheet metal, cardboard or similar rigid tubing securely attached to rafters or other horizontal surfaces where rats will pass. Employ tubes long enough to minimize spillage of powder through the ends.

APPLICATION DIRECTIONS: For House mice use at the rate of 1 oz. powder per 2.5 sq. ft. of runway area. For Norway and Roof rats use at the rate of 2 oz. powder for 2.5 sq. ft. of runway area. Apply the powder into the rat burrow or holes or within walls with a hand bulb or similar duster. Do not use power dusting devices. Sprinkle the powder in patches in such a manner as to expose the rats and house mice to it. Patch size may be (but not limited to) 6 inches x 12 inches and should be adapted to each situation. For burrows that are located along the periphery of buildings and that are likely to serve as routes of entry into these structures, place about 5 grams of tracking powder in each burrow. Close burrows with soil, loose leaves or paper. Repeat treatment if burrows are reopened. Maintain powder in treated areas for at least 20 days. Wearing gloves, collect and dispose of all visible dead animals and used powder promptly. Repeat treatment when infestation recurs. Where a continuous source of infestation is present, establish permanent tracking powder stations and replenish the powder as needed. (083011)